VOL. 750, NOS. 1 + 2 JOURNAL OF CHROMATOGRAPHY A 25 OCTOBER 1996

CONTENTS

(Abstracts/Contents Lists published in Analytical Abstracts, Biochemical Abstracts, Biological Abstracts, Chemical Abstracts, Chemical Titles, Chromatography Abstracts, Current Awareness in Biological Sciences (CABS), Current Contents/Life Sciences, Current Contents/Physical, Chemical & Earth Sciences, Deep-Sea Research/Part B: Oceanographic Literature Review, Excerpta Medica, Index Medicus, Mass Spectrometry Bulletin, PASCAL-CNRS, Referativnyi Zhurnal, Research Alert and Science Citation Index)

4TH INTERNATIONAL SYMPOSIUM ON HYPHENATED TECHNIQUES IN CHROMATOGRAPHY AND HYPHENATED CHROMATOGRAPHIC ANALYZERS, BRUGES, 7–9 FEBRUARY 1996

Preface
by R. Smits (Antwerp, Belgium) 1

REVIEWS

Future potential of targeted component analysis by multidimensional liquid chromatography–mass spectrometry
by F. Regnier and G. Huang (West Lafayette, IN, USA) 3

Vaporising systems for large volume injection or on-line transfer into gas chromatography: classification, critical remarks and suggestions
by K. Grob and M. Biedermann (Zürich, Switzerland) 11

Improvement of quality control of speciation analysis using hyphenated techniques. A decade of progress within the European Community
by Ph. Quevauviller (Brussels, Belgium) 25

SYSTEMS INVOLVING MS COUPLING

Application of multidimensional affinity high-performance liquid chromatography and electrospray ionization liquid chromatography–mass spectrometry to the characterization of glycosylation in single-chain plasminogen activator. Initial results
by A. Apffel, J.A. Chakel and W.S. Hancock (Palo Alto, CA, USA) and C. Souders, T. M'Timkulu and E. Pungor, Jr. (Brisbane, CA, USA) 35

Gas and liquid chromatography coupled to tandem mass spectrometry for the multiresidue analysis of β-agonists in biological matrices
by G. Van Vyncht (Liège, Belgium), S. Preece (Altrincham, UK) and P. Gaspar, G. Maghuin-Rogister and E. DePauw (Liège, Belgium) 43

Hyphenation of capillary electrophoresis to inductively coupled plasma mass spectrometry as an element-specific detection method for metal specification
by B. Michalke and P. Schramel (Oberschleißheim, Germany) 51

What is the true odour of cut *Allium*? Complementarity of various hyphenated methods: gas chromatography–mass spectrometry and high-performance liquid chromatography–mass spectrometry with particle beam and atmospheric pressure ionization interfaces in sulphenic acids rearrangement components discrimination
by S. Ferary and J. Auger (Tours, France) 63

Quantitative determination of norethisterone acetate in human plasma by capillary gas chromatography with mass-selective detection
by F. Pommier, A. Sioufi and J. Godbillon (Rueil Malmaison, France) 75

Simultaneous determination of arsenic species by ion chromatography–inductively coupled plasma mass spectrometry
by P. Teräsahde (Helsinki, Finland) and M. Pantsar-Kallio and P.K.G. Manninen (Lahti, Finland) 83

Speciation of chromium in waste waters by coupled column ion chromatography–inductively coupled plasma mass spectrometry
by M. Pantsar-Kallio and P.K.G. Manninen (Lahti, Finland) 89

Coupled simulated distillation–mass spectrometry for the evaluation of hydroconverted petroleum residues
by R. Bacaud and L. Rouleau (Villeurbanne, France) 97

Comparison of the possibilities of gas chromatography–mass spectrometry and tandem mass spectrometry systems for the analysis of anabolics in biological material
by H.F. De Brabander and P. Batjoens (Merelbeke, Belgium), D. Courtheyn and J. Vercammen (Gentbrugge, Belgium) and K. De Wasch (Merelbeke, Belgium) 105

On-line liquid chromatography–electrospray ionization mass spectrometry for determination of the brevetoxin profile in natural "red tide" algae blooms
by Y. Hua and W. Lu (New Orleans, LA, USA), M.S. Henry and R.H. Pierce (Sarasota, FL, USA) and R.B. Cole (New Orleans, LA, USA) 115

Rapid and high-performance analysis of thyreostatic drug residues in urine using gas chromatography–mass spectrometry
by P. Batjoens, H.F. De Brabander and K. De Wasch (Merelbeke, Belgium) 127

Gas chromatographic–tandem mass spectrometric analysis of clenbuterol residues in faeces
by P. Batjoens (Merelbeke, Belgium), D. Courtheyn (Gentbrugge, Belgium), H.F. De Brabander (Merelbeke, Belgium), J. Vercammen (Gentbrugge, Belgium), K. De Wasch (Merelbeke, Belgium) and M. Logghe (Gentbrugge, Belgium) 133

On-line monitoring of biotechnological processes by gas chromatographic–mass spectrometric analysis of fermentation suspensions
by G. Matz and F. Lennemann (Hamburg, Germany) 141

A new fast method for field screening of polychlorinated biphenyls in air
by G. Matz, W. Schröder and T. Ollesch (Hamburg, Germany) 151

Pyrolytic alkylation–gas chromatography–mass spectrometry of model polymers. Further insights into the mechanism and scope of the technique
by F.J. González-Vila, J.C. Del Rio, F. Martin and T. Verdejo (Seville, Spain) 155

Automated at-line solid-phase extraction–gas chromatographic analysis of micropollutants in water using the PrepStation
by Th. Hankemeier, P.C. Steketee, J.J. Vreuls and U.A.Th. Brinkman (Amsterdam, Netherlands) 161

On-line coupled supercritical fluid extraction and chromatography for the determination of thiolcarbamate herbicides in soil matrix
by T. Yarita, A. Nomura, Y. Horimoto and S. Gonda (Ibaraki, Japan) 175

Analysis of poly(vinyl chloride) additives by supercritical fluid extraction and gas chromatography
by M.L. Marín, A. Jiménez, J. López and J. Vilaplana (Alicante, Spain) 183

Optimization of supercritical fluid extraction–gas chromatography of methylmercury in marine samples
by R. Cela-Torrijos, M. Miguéns-Rodríguez, A.M. Carro-Díaz and R.A. Lorenzo-Ferreira (Santiago de Compostela, Spain) 191

Fat content for nutritional labeling by supercritical fluid extraction and an on-line lipase catalyzed reaction
by J.M. Snyder, J.W. King and M.A. Jackson (Peoria, IL, USA) 201

Automated on-line solid-phase extraction–high-performance liquid chromatography–diode array detection of phenolic compounds in sherry wine
by C. Chilla, D.A. Guillén, C.G. Barroso and J.A. Pérez-Bustamante (Cádiz, Spain) 209

Investigation of parameters affecting the on-line combination of supercritical fluid extraction with capillary gas chromatography
by X. Lou, H.-G. Janssen and C.A. Cramers (Eindhoven, Netherlands) 215

Solid-phase extraction of polar pesticides from environmental water samples on graphitized carbon and Empore-activated carbon disks and on-line coupling to octadecyl-bonded silica analytical columns
by J. Slobodník, Ö. Öztezkizan, H. Lingeman and U.A.Th. Brinkman (Amsterdam, Netherlands) 227

Coupling of thermal desorption in modified closeable sampling columns with wide-bore capillary gas chromatography and mass spectrometric detection (Short communication)
by H.G. Struppe, F. Franke, J. Hofmann and B. Ondruschka (Leipzig, Germany) 239

Application of ^{13}C-labelled polychlorinated biphenyl congener 153 as internal standard in the gas chromatographic–mass spectrometric analysis of polychlorinated biphenyls (Short communication)
by A. Kubátová and M. Matucha (Prague, Czech Republic) and M. Bubner (Rossendorf, Germany) 245

SYSTEMS WITH DIODE ARRAY DETECTION

Determination of taxol in *Taxus* species grown in Hungary by high-performance liquid chromatography–diode array detection. Effect of vegetative period
by V. Németh-Kiss, E. Forgács, T. Cserháti and G. Schmidt (Budapest, Hungary) 253

Simultaneous determination of 3,4-dimethoxybenzaldehyde and 3,4-dimethoxyphenylacetone in industrial waste waters by high-performance liquid chromatography–diode array detection
by T. Cserháti and E. Forgács (Budapest, Hungary) 257

LC–LC AND LC–GC COUPLING

Determination of polycyclic aromatic hydrocarbons in edible oils and fats by on-line donor–acceptor complex chromatography and high-performance liquid chromatography with fluorescence detection
by F. van Stijn, M.A.T. Kerkhoff and B.G.M. Vandeginste (Vlaardingen, Netherlands) 263

Automated on-line gel permeation chromatography–gas chromatography for the determination of organophosphorus pesticides in olive oil
by J.J. Vreuls, R.J.J. Swen and V.P. Goudriaan (Amsterdam, Netherlands), M.A.T. Kerkhoff and G.A. Jongenotter (Vlaardingen, Netherlands) and U.A.Th. Brinkman (Amsterdam, Netherlands) 275

OTHER TYPES OF HYPHENATED AND MULTIPLE HYPHENATION PROCEDURES

Hyphenated techniques for characterizing coal wastewaters and associated sediments
by J. Pörschmann, F.-D. Kopinke, M. Remmler, K. Mackenzie, W. Geyer and S. Mothes (Leipzig, Germany) . . . 287

Application of on-line capillary high-performance liquid chromatography–nuclear magnetic resonance spectrometry coupling for the analysis of vitamin A derivatives
by K. Albert, G. Schlotterbeck and L.-H. Tseng (Tübingen, Germany) and U. Braumann (Rheinstetten, Germany) 303

Identification of disaccharides by gas chromatography–Fourier transform infrared spectroscopy
by R.G. Veness and C.S. Evans (London, UK) 311

Determination of butyl- and phenyltin compounds in biological material by gas chromatography–flame photometric detection after ethylation with sodium tetraethylborate
by C. Carlier-Pinasseau, A. Astruc, G. Lespes and M. Astruc (Pau, France) 317

Optimization of the determination of inorganic and organic selenium species using high-performance liquid chromatography–electrothermal atomic absorption spectrometry
by N. Gilon, M. Potin-Gautier and M. Astruc (Pau, France) 327

Surface ionization detector with a supersonic free jet for gas chromatography. Some applications
by H. Kishi (Oyama, Japan) and T. Fujii and G. Sato (Ibaraki, Japan) 335

Quantification of two chromatographic unresolved dichlorophenols using gas chromatography–direct deposition-Fourier transform infrared spectrometry and multivariate calibration
by I. Rodríguez, M.H. Bollaín and R. Cela (Compostela, Spain) 341

Automated determination of verapamil and norverapamil in human plasma with on-line coupling of dialysis to high-performance liquid chromatography and fluorometric detection
by A. Ceccato, P. Chiap, Ph. Hubert, B. Toussaint and J. Crommen (Liège, Belgium) 351

On-line high-performance liquid chromatography–solvent evaporation–high-performance liquid chromatography–capillary gas chromatography–flame ionisation detection for the analysis of mineral oil polyaromatic hydrocarbons in fatty foods
by S. Moret and K. Grob (Zürich, Switzerland) and L.S. Conte (Udine, Italy) 361

Pesticide residue analysis in foodstuffs applying capillary gas chromatography with atomic emission detection. State-of-the-art use of modified multimethod S19 of the Deutsche Forschungsgemeinschaft and automated large-volume injection with programmed-temperature vaporization and solvent venting
by H.-J. Stan and M. Linkerhägner (Berlin, Germany) 369

Extraction of maleic hydrazide residues from potato crisps and their determination using high-performance liquid chromatography with UV and atmospheric pressure chemical ionisation mass spectrometric detection (Short communication)
by D.J. Lewis, K.A. Barnes, K. Wilkinson, S.A. Thorpe, S.L. Reynolds and J.R. Startin (Norwich, UK) 391

Quantification of cyclamate and cyclohexylamine in urine samples using high-performance liquid chromatography with trinitrobenzenesulfonic acid pre-column derivatization (Short communication)
by I. Casals, M. Reixach, J. Amat, M. Fuentes and L. Serra-Majem (Barcelona, Spain) 397

Toxicological evaluation of harmful substances by in situ enzymatic and biological detection in high-performance thin-layer chromatography (Short communication)
by C. Weins and H. Jork (Saarbrücken, Germany) 403

AUTHOR INDEX 409